

# The Top 10 Things You Need to Know About Acquired Pigmentation Disorders

Maggie L. Chow, Janiene Luke, Sharon E. Jacob

**ABSTRACT:** Disorders of pigmentation can be acquired or inherited. The acquired disorders present an opportunity for surveillance and treatment to prevent morbidity and improve quality of life. In this article, we delineate the acquired causes, pathogenesis, and treatment options of disorders of skin hyperpigmentation or hypopigmentation. **Key words:** Acquired Pigmentation Disorders, Dyspigmentation, Hyperpigmentation, Pityriasis alba, Vitiligo

**M**elanin, a brown or black pigment, determines the color of skin and hair. This pigment is formed in melanocytes from tyrosine (Hurwitz, 2006). Melanocytes are cells derived from the neural crest that migrate to the basal layer of the epidermis during embryogenesis and become dendritic secretory cells when mature. Abnormal skin hyperpigmentation or hypopigmentation results from aberrant melanin production, melanosome transport, or melanocyte development.

Defects in the melanin production pathway can be inherited or acquired. The genodermatoses related to disorders of pigmentation are described in a separate article (Chow &

Jacob, 2015). Here, we describe the “top 10 things to know about” the disorders of hyperpigmentation and hypopigmentation caused by functional abnormalities secondary to an acquired extrinsic or intrinsic cause.

## 1. Acquired Versus Inherited Pigmentary Disorders

Acquired disorders are medical conditions that develop over the course of an individual’s life. In contrast, an inherited pigmentary disorder arises secondarily to genes passed on from parents to their offspring. Thus, an acquired pigmentary disorder is hypopigmentation or hyperpigmentation that is not present at birth and is not secondary to an inherited genetic defect from the parents. Examples of such disorders include vitiligo, melasma, and postinflammatory pigment alterations secondary to tinea versicolor, among others. In contrast, inherited pigmentary disorders are secondary to a genetic aberrancy and can include such syndromes as tuberous sclerosis, neurofibromatosis, and the Carney complex.

## 2. Pathophysiology

Human skin color is primarily determined by the type, quantity, and distribution of melanin pigment in the skin, which is regulated by a complex array of genetic, environmental, and endocrine factors (Costin & Hearing, 2007). Aside from determining skin color, melanin also protects the skin against ultraviolet (UV) radiation. Extrinsic and intrinsic disturbances in the physiologic and biological processes that affect the production and transport of melanin, such as UV light, certain medications, and the hormonal changes that occur during pregnancy, can also affect skin color.

Melanocytes are the secretory cells that produce melanin. The quantity of melanocytes in human skin across ethnicities is constant within each individual; however, skin color is determined by the size, number, and distribution of melanosomes within keratinocytes (Haake & Holbrook, 1999). Melanocytes reside in the basal layer of the epidermis. After the production of melanin, the melanocytes transfer

Maggie L. Chow, PhD, MD, Kaiser Permanente Los Angeles Medical Center, Los Angeles, CA.

Janiene Luke, MD, Loma Linda University, Loma Linda, CA.

Sharon E. Jacob, MD, Loma Linda University, Loma Linda, CA.

Sharon E. Jacob served as an independent investigator on the safety and efficacy of T.R.U.E. Test (Smart Practice, Phoenix, AZ) Panels 1.1, 2.1, and 3.1 in children and adolescents, Pediatric Research Equity Act (PREA-1) trial, and now serves as an investigator on PREA-2. She has served as a consultant for Johnson & Johnson. She has no conflicts of interest associated with the subject matter in this manuscript. Dr. Luke has no relevant disclosures or conflicts of interest. Dr. Chow has no relevant disclosures or conflicts of interest.

Correspondence concerning this article should be addressed to Sharon E. Jacob, MD, Department of Dermatology, Loma Linda University, 11370 Anderson Street, Suite 2600, Loma Linda, CA 92354.

E-mail: sjacob@contactderm.net

DOI: 10.1097/JDN.0000000000000169

the pigment to surrounding keratinocytes through melanosomes, which are specialized membrane-bound organelles (Costin & Hearing, 2007).

The synthesis and distribution of melanin depends on several steps. First, melanoblasts (melanocyte precursor cells) must develop and migrate from the neural crest to their final peripheral sites. Second, the melanoblasts must differentiate into melanocytes after migration. Finally, the melanocytes must then proliferate and begin producing melanosomes, which must undergo a maturation process. Three enzymes are required to synthesize melanin within these organelles, including tyrosinase, dopachrome tautomerase, and tyrosinase-related protein 1 (Haake & Holbrook, 1999).

Melanosomes can produce several types of melanins: eumelanin (black-brown), pheomelanin (yellow-red), a combination of eumelanin and pheomelanin termed mixed melanins, and neuromelanin. These pigments differ in their chemical, structural, and physical properties and play disparate molecular roles in different anatomic regions (Hearing, 2000). Once the melanosomes are transferred to neighboring keratinocytes, the epidermal melanin unit becomes a functional entity that can respond through paracrine and autocrine mechanisms to environmental stimuli. Thus, human skin pigmentation is a dynamic process determined by numerous internal and external factors (Hearing, 2000).

### **3. Causes of Human Skin Hypopigmentation or Hyperpigmentation**

Internal and external processes that affect the steps in the complex melanin production and transport pathway can potentially lead to abnormalities in skin pigmentation. Internal processes include autoimmune etiologies (e.g., vitiligo) and endocrine dysfunction (e.g., insulin resistance). In contrast, external processes can include medications (e.g., minocycline), sun exposure (e.g., solar lentigo), and post-inflammatory hyperpigmentation responses to injury and infections (e.g., tinea versicolor).

For instance, numerous medications can cause skin discoloration. Heavy metals, such as silver, bismuth, arsenic, and gold, can cause hyperpigmentation through binding and inactivating sulfhydryl compounds that function to inhibit tyrosinase activity. This process results in melanogenesis (Molokhia & Portnoy, 1973). Chemotherapy agents are hypothesized to cause hyperpigmentation through toxic effects on melanocytes, stimulation of melanocytes, and inflammation (Costin & Hearing, 2007). Levodopa, which is used for the treatment of Parkinson's disease, is hypothesized to enhance melanin biosynthesis through possible extracellular oxidation (Hendrix & Greer, 1992). Other medications that have been associated with skin hyperpigmentation include sulfonamide and tetracycline antibiotics, nonsteroidal anti-inflammatory medications, diuretics, hydantoin antiepileptics, and psychoactive medications (Levantine & Almeyda, 1973). The pathophysiology of common causes of hypopigmentation and hyperpigmentation are described in Tables 1 and 2.

### **4. Epidemiology and Risk Factors**

The epidemiology of acquired pigmentation disorders varies widely by disease association. These are described in Tables 1 and 2. Dyspigmentation can affect the young or the older adults and both men and women around the world. These epidemiologic clues can aid in diagnosis. For example, tinea versicolor should be near the top of the differential diagnosis for scaly hypopigmented macules on the chest of a teenaged patient living in a humid environment. In endemic regions, infectious causes must also be ruled out; for example, pinta is a treponemal disease that is similar to syphilis and causes hypopigmentation.

### **5. Prevention**

A critical intervention in the prevention of hyperpigmentation is sun protection, as exposure can exacerbate the dark areas through deposition of melanin into the skin. Sun exposure can lead to both acute and chronic pigmentary darkening. In the immediate (acute) setting, UV light (specifically UVA and, to a lesser extent, UVB) oxidizes tyrosine in the skin and ultimately forms melanin, causing skin darkening (Joshi, Carraro, & Pathak, 1987). In the chronic setting, hyperpigmentation secondary to UVA/UVB exposure arises from activation and proliferation of the exposed melanocytes (Gilcrest, Blog, & Szabo, 1979).

Furthermore, the cessation of aggravating or inciting exogenous factors should be pursued, and infections should be treated appropriately. Underlying dermatoses, such as acne, must be treated, and allergens and irritants need to be avoided in contact dermatitis to prevent further development of dyspigmented lesions.

### **6. The Social Impact of Dyspigmentation**

Patients with dyspigmentation, for example, vitiligo and pigmented contact dermatitis, both of which can affect the face, may have their quality of life significantly affected secondary to the impact that their disease has on appearance. Patients may experience depression, anger, embarrassment, and shame. They may worry that their disease will continue to progress (Sampogna et al., 2008) or be contagious. Facial lesions in particular can be associated with low self-esteem, social withdrawal, and lower school and work productivity. Furthermore, patients who elect to pursue medical treatments and procedures can experience high financial burdens for results that may not reach their expectations (Handel, Miot, & Miot, 2014).

Thus, it is important to discuss the etiology of the dyspigmentation with patients and explore their level of distress involving their disorder. Patients should be reassured of their prognosis but also given realistic expectations regarding the efficacy of their treatments.

### **7. Treatment**

Treatment options for hypopigmentation or hyperpigmentation include medical and surgical options. In either

**TABLE 1. Hyperpigmented Lesions**

Lesion Name	Presentation	At-Risk Populations/Epidemiology	Pathophysiology	Treatment Options
Postinflammatory hyperpigmentation	Hyperpigmented, irregularly shaped macules and patches in an area after previous inflammation or injury	More common in darker skin types	The inflammatory response causes arachidonic acid to be released and oxidized into such products as leukotrienes and prostaglandins, which stimulate epidermal melanocytes. This stimulation causes increased production of melanin, which is then transferred to surrounding keratinocytes, resulting in epidermal hypermelanosis.	Daily sun protection (SPF > 30) Hydroquinone 2%–12% Tretinoin cream 0.05% ointment (Reti-A) All-trans retinoic acid aqueous gel 0.1%–0.4% with hydroquinone-lactic acid ointment Salicylic 0.5%–2% or glycolic acid 10% peels Azelaic acid 20% cream (Azelex) Tazarotene 0.1% cream Aloe vera leaf extract Glabridin microsponge-loaded gel
Melasma	Tan-to-brown progressive skin hyperpigmentation usually of the face that can occur in malar, centofacial, and mandibular patterns and less frequently isolated on the forearms	Affects over 5 million people in the United States (Grimes, 1995); Prevalence is 8.8%–40% (Werlinger et al., 2007); More common in women > men, pregnant women (15%–50%), women of childbearing age, family history of melasma, and light-brown skin types (Latinos and Asians)	Dermal hyperpigmentation is caused by ‘‘pigmentary incontinence,’’ when pigment is trapped by macrophages in the papillary dermis after inflammation-related damage to the basal cell layer (Cardinali, Kovacs, & Picardo, 2012).  The factors associated with the development of melasma include genetic disposition, sunlight exposure, and hormonal influence (Hughes, 1987; Ortonne et al., 2009). However, the exact mechanism by which melasma arises remains unclear.	Aggressive sun protection (SPF > 30), and cessation of any exposure to estrogen. Hydroquinone Tretinoin, adapalene, and azelaic acid Combination of bleaching agents, for example, Tri-Luma, which is composed of tretinoin 0.05%, hydroquinone 4%, and fluocinolone acetonide 0.01% (Baliña & Graupe, 1991; Dogra, Kanwar, & Paisad, 2002; Grimes et al., 2010; Sheih & Pandya, 2011). Chemical peels can help a subset of patients (Rivas & Pandya, 2013; Taylor & Anderson, 1994). Laser therapy (judicious)

(continues)

**TABLE 1. Hyperpigmented Lesions, Continued**

<b>Lesion Name</b>	<b>Presentation</b>	<b>At-Risk Populations/Epidemiology</b>	<b>Pathophysiology</b>	<b>Treatment Options</b>
Solar lentiginos	Sharply demarcated hyperpigmented macular lesions that increase in size over time and can appear on the hands, face, neck, and forearms (Ortonne, Pandya, Lui, & Hexsel, 2006)	Patients with history of sun exposure	May be caused by repeated past UV light exposure leading to mutagenesis and increased melanin production; Can also serve as a marker for previous sun burns	Cryotherapy (most common and efficacious) Bleaching agents (including 2% mequinol and 0.01% tretinoin) Tri-Luma Over-the-counter skin-lightening agents targeting the melanin production pathway
Erythema dyschromicum perstans (Ashy dermatoses)	Hyperpigmented blue-gray oval or round patches on the face, trunk, and extremities	Most common in Latin America and Asia; More common in darker skinned individuals, women more commonly than men (Zaynoun, Rubeiz, & Kibbi, 2008)	Hypothesized to be mediated by an immunologic mechanism (Gillbro & Olsson, 2011); May also be caused by chloroethanol, ammonium nitrate ingestion, x-ray contrast media ingestion, other environmental contaminants, and chronic Hepatitis C infection	Spontaneous remission Narrow-band UVB Dapsone Clotazimine (Gillbro & Olsson, 2011; Torreló et al., 2005)
Pigmented contact dermatitis (Riehl melanosis; Serrano, Pujol, Cuadra, Gallo, & Allaga, 1989)	Brownish gray hyperpigmented patches in a reticular pattern with or without erythematous macules or papules that can be especially prominent on the forehead and temples but can be present on the dorsal hands, forearms, and neck	Usually occurs in young and middle-aged women; epidemiological studies not conducted	Contact allergy, most commonly to cosmetic chemicals	Patch testing Removal of causative agent Sun protection Hydroquinone
Poikiloderma of Civatte	Symmetric red-brown hyperpigmented patches in a reticulate pattern	Middle-aged or elderly fair-skinned women; Epidemiology unknown	Chronic sun exposure and hormonal factors, especially in individuals with genetic susceptibility to solar radiation (Katoulis et al., 1999); Photosensitization by chemicals in cosmetics (Katoulis et al., 2002)	Ablative fractional laser resurfacing Pulse dye laser (Tierney & Hanke, 2009)

(continues)

**TABLE 1. Hyperpigmented Lesions, Continued**

Lesion Name	Presentation	At-Risk Populations/Epidemiology	Pathophysiology	Treatment Options
Drug-induced hyperpigmentation	Slow onset and progression of brownish to blue-gray discoloration favoring sun-exposed areas	Patients taking tricyclic antidepressants, anticonvulsants, antimalarial medications, amiodarone, cytotoxic medications, minocycline, and phenothiazines; Accounts for 10%–20% of all acquired hyperpigmentation worldwide. Dark skin > fair skin (Haider, Nandedkar, & Neal, 2003)	Stimulation of epidermal melanocytes leading to melanin hyperproduction, impairment of dermal macrophage melanin clearance, induction of special pigment production, accumulation of the medication in the skin, and vessel damage leading to iron deposition (Dereure, 2001)	Discontinuation of the medication

The presentation, pathophysiology, and treatment options for acquired hyperpigmented lesions are listed.

case, aggressive sun protection (at least SPF 30) is the cornerstone of any therapeutic regimen. In general, the medical treatments available for skin hyperpigmentation include measures to prevent underlying dermatoses causing hyperpigmentation (e.g., acne) and to remove offending agents (e.g., long-term minocycline use). Topical retinoids and bleaching agents have varying efficacy depending on the etiology of the condition leading to the pigment alteration. Although topical hydroquinone (both prescription strength and over the counter) has become a standard therapy for hyperpigmentation, current evidence supports combination therapy utilizing several modalities. For example, triple combination therapy with fluocinolone acetonide 0.01%, hydroquinone 4%, and tretinoin 0.05% given for 8 weeks has been associated with greater improvement of melasma by standardized measurements compared with hydroquinone standard therapy (Chan et al., 2008; see Tables 1 and 2).

Several laser and surgical treatments are also available for hyperpigmentation. Laser therapies, including pulse dye laser and ablative fractional laser resurfacing, have shown some efficacy for hyperpigmented patches, particularly for lentigines and postinflammatory hyperpigmentation (Ho et al., 2012; Ong & Bashir, 2012). That said, there may be a significant risk of postlaser pigment alteration, and thus, it is important to first assess for untoward laser effects by performing a test spot in an inconspicuous region, for example, behind the ear. In extreme cases, surgical excision of the dyspigmented region followed by skin grafting might be pursued (Al-Qattan, 2000).

The treatments for hypopigmentation vary based on the mechanism of dyspigmentation and also include medical and surgical treatments. For example, in the autoimmune disorder of vitiligo, ruling out underlying disorders, such as thyroid disease, is necessary. Once this has been done, the patient could opt for treatment with a topical corticosteroid or an immunomodulatory topical calcineurin inhibitor. Infections should be treated with appropriate antifungal or antibacterial therapies. In select cases, phototherapy, excimer laser, and graft surgery may play a role in treatment.

Narrow-band light therapy can also be helpful in some cases of acquired hypopigmentation or depigmentation. As a general principle, no singular therapy is completely efficacious, and combination therapies have been shown to yield the best results.

### 8. Follow-up

Follow-up after the initiation of treatment depends on the proposed regimen and the need for surveillance. In general, patients with dyspigmentation can be reassessed every 3–6 months for evaluation of treatment efficacy and adjustment of the regimen. This also assures that patients are appropriately using their medications and that physicians can withdraw the medications at the earliest sign of adverse event (e.g., exogenous ochronosis secondary to hydroquinone).



**TABLE 2. Hypopigmented Lesions**

Lesion Name	Presentation	At-Risk Populations/ Epidemiology	Pathophysiology	Treatment Options
Vitiligo (Taieb et al., 2013)	Hypopigmented or milk-white sharply demarcated macules and patches that enlarge centrifugally over time; can be localized, generalized, or universal	Individuals with other autoimmune disease, such as Graves' disease, Addison, diabetes mellitus, alopecia areata, pernicious anemia, inflammatory bowel disease, and psoriasis. Relative rate of 1%–2%. Thirty percent of cases will have family history of vitiligo (Ortonne, 2008).	Etiology is multifactorial. Leading theories include autoimmune or cytotoxic destruction of melanocytes and intrinsic melanocyte defects. Oxidative stress and neural injury may play a role in some subtypes of vitiligo.	Topical corticosteroids Pimecrolimus and tacrolimus Phototherapy Excimer laser Graft surgery
Pityriasis alba (Lin & Janniger, 2005)	Multiple symmetric, poorly defined macules and patches with pityriasisiform scaling over the face	Children under 16 years old; Prevalence in children of 5% (Kim & Lockey, 2010)	Reduction in active melanocytes and decreased melanosomes and melanosome size in affected skin	Emollients Low-potency topical corticosteroids for erythema or pruritus Excimer laser (Al-Mutairi & Hadad, 2012; Gambichler et al., 2007; Rigopoulos et al., 2006) Topical calcineurin inhibitors
Tinea versicolor (Crespo-Erchiga & Florencio, 2006)	Hypopigmented (can also be hyperpigmented) macules and patches on the back or chest	Residents of humid and hot climates, most common ages of 15–24 years; Prevalence of 2%–8% in the national population (Nehoff, Krüger, Ginter-Hanselmayer, & Tietz, 2014)	Infection by <i>Malassezia globosa</i> and <i>Malassezia furfur</i> in the stratum corneum. Genetic predisposition, immunosuppression, Cushing disease, malnutrition, and warm, humid environments	Topical selenium sulfide Topical ciclopirox Topical antifungal medications Topical sodium sulfacetamide Oral fluconazole or itraconazole
Postinflammatory hypopigmentation (Nicolaidou & Katsambas, 2014)	Hypopigmented macules and patches in an area of previous injury or inflammation	Patients with atopic dermatitis, psoriasis, seborrheic dermatitis, sarcoidosis, lupus, or mycoses fungoides	Effect of inflammation and injury on the melanin production and transportation pathway	Spontaneous resolution UVB phototherapy/ excimer laser Epidermal or melanocyte grafting
Hansen's Disease (Leprosy; Britton & Lockwood, 2004; Ustianowski & Lockwood, 2003)	Painless, insensate, and possibly hypopigmented patch	Travelers to or individuals from endemic tropical regions; In the United States, 200–300 cases of leprosy reported each year particularly in states with large immigrant populations; also endemic in Texas, Louisiana, and Hawaii	Infection with <i>M. leprae</i>	For paucibacillary leprosy, dapsone 100 mg self-administered daily plus rifampicin 600 mg monthly supervised for 6–12 months

(continues)

**TABLE 2. Hypopigmented Lesions, Continued**

Lesion Name	Presentation	At-Risk Populations/ Epidemiology	Pathophysiology	Treatment Options
		Lepromatous type 2:1 in men > women; Bimodal age distribution ("Global leprosy update, 2013; reducing disease burden," 2014)		For multibacillary leprosy, dapsons 100 mg self-administered daily plus rifampicin 600 mg monthly supervised for 24 months or clofazimine 50 mg self-administered daily plus clofazimine 300 mg monthly supervised for 24 months  In cases with multisystem involvement, consultation with ophthalmology, plastic surgery, neurosurgery, orthopedic surgery, and otolaryngology may be necessary.

The presentation, pathophysiology, and treatment options for acquired hypopigmented lesions are listed.

### 9. The Challenges

An important point to address in the treatment of dyspigmentation is that no single treatment is 100% effective. Combination therapies, both surgical and medical, often yield the best results. Furthermore, the treatments themselves may lead to hypopigmentation or depigmentation at previous lesions, and the treatment of hypopigmentation can lead to hyperpigmentation at previous lesions. For example, hydroquinone has been reported as a cause of secondary hyperpigmentation (exogenous ochronosis) when used for extended periods. Fortunately, this side effect is extremely rare (Makino et al., 2013).

Another important issue to consider is the ethics involving treatments for dyspigmentation (Baumann, 2012), as many dermatology practices now are selling products for skin lightening. This issue has been hotly debated among dermatologists for decades and continues to be a topic for discussion. The American Academy of Dermatology position statement on in-office dispensing includes only prescribing medications in the best interest of the patients in an evidence-based manner (American Academy of Dermatology, 1999). It has been suggested that guidelines delineating an objective, organized approach for therapy be created to lead practitioners and their patients in treatment.

Recently, cosmetic products used to treat dyspigmentation have been a cause for concern. In the United States, it is illegal to use mercury as an ingredient in skin bleaching products. However, skin-lightening products from other countries have been noted to contain this potentially hazardous substance (Boonchai, Desomchoke, & Iamtharachai, 2011; Cristaudo et al., 2013). Mercury is an efficacious

active ingredient because it competes with copper in the tyrosinase-regulated step of the melanin synthesis pathway, thereby decreasing pigment production (Vazirnia & Jacob, 2014). However, the metal is an emerging cause of skin hypersensitivity (Forte, Petrucci, & Bocca, 2008) and an increasingly prevalent cause of metal-induced allergic contact dermatitis. Thus, it is important to warn patients about the potential risks of sensitization when using cosmetic products from outside the United States for dyspigmentation.

### 10. On the Horizon

The treatment of dyspigmentation is an active field of study, with exciting new and emerging clinical studies identifying more and more efficacious treatment modalities for this potentially morbid condition. Moreover, because patients with ethnic skin may experience increased morbidity from dyspigmentation, ongoing research on treatments specifically for this population are extremely important (Alexis, 2013).

Recent data suggest that retinoids and azelaic acid are highly efficacious for treatment of both acne and postinflammatory hyperpigmentation (Woolery-Lloyd, Keri, & Doig, 2013). Other newly developed formulations have also shown efficacy for treating hyperpigmentation. A 12-week randomized, double-blinded, half-face study revealed greater patient satisfaction and improvement in hyperpigmentation in subjects receiving a brightener formulation containing SMA-432 (a prostaglandin E2 inhibitor) over 4% hydroquinone. Encouragingly, 95% of subjects reported improvement in overall skin condition at the end of the study (Makino et al., 2013). Other medical treatments on the horizon for hyperpigmentation treatment include topical methimazole

(peroxidase inhibitor), 4-n-butylresorcinol (tyrosinase inhibitor), zinc, arbutin (Makino et al., 2013), soy protein, N-acetyl glucosamine, kojic acid, aleosin, licorice extract, and ascorbic acid (Draelos, 2007).

The use of laser treatment in combination with medical therapies for hyperpigmentation is also an active field of research, although the efficacy of this treatment has been variable (Arora, Sarkar, Garg, & Arya, 2012; Vashi & Kundu, 2013). Recently, a study examining the efficacy of low-fluence Q-switched Nd:YAG laser treatment immediately after microdermabrasion in combination with hydroquinone and tretinoin or vitamin C showed >50% clearance of melasma after 1 month (Kauvar, 2012). The 1927-nm thulium fiber fractional laser has also been shown to improve melasma histologically in a split-face study of 25 patients (Lee, Haw, Kim, Chang, & Lee, 2013). Other studies, however, have yielded mixed conclusions (for a review, please see Arora et al., 2012). Despite these results, the interpretation of our current studies is limited by the short duration of follow-up, and longer-term studies are needed to assess long-term treatment efficacy and adverse effects. It is our experience that melasma often recurs or may even be exacerbated after laser treatment, and thus, caution must be exercised when considering these treatments. New therapies like these continue to emerge, allowing dermatologic providers a wider range of management options for hyperpigmentation.

The emerging therapies for hypopigmentation are mainly focused on vitiligo, given that we have efficacious treatments for the infectious causes of hypopigmentation. Recently, the HI-Light trial showed that lightweight handheld NB-UVB units for the treatment of vitiligo were associated with good treatment adherence and repigmentation rates (Eleftheriadou et al., 2014). Among the surgical methods for vitiligo treatment, split-thickness skin grafts had the highest repigmentation success rates over punch/mini-graft, cultured and noncultured cellular transplantation, and blister roof grafting (Mulekar & Isedeh, 2013). These approaches are among those being investigated, although standardized head-to-head trials for the varied treatment methods are needed (Eleftheriadou, 2013).

## SUMMARY

In summary, the acquired pigmentary disorders are a heterogeneous set of conditions that can cause severe distress and quality-of-life issues. In contrast to the inherited pigmentary disorders, these conditions present an opportunity for prevention and early intervention. The treatment of dyspigmentation remains an art and must be individually tailored to each patient. Our understanding about the pathophysiology of each disorder continues to improve, but long-term studies on treatment efficacy and adverse effects are needed to test the many proposed therapies. ■

## REFERENCES

Alexis, A. F. (2013). Lasers and light-based therapies in ethnic skin: Treatment options and recommendations for Fitzpatrick skin types V and VI. *The British Journal of Dermatology*, 169(Suppl. 3), 91–97. doi:10.1111/bjd.12526

- Al-Mutairi, N., & Hadad, A. A. (2012). Efficacy of 308-nm xenon chloride excimer laser in pityriasis alba. *Dermatologic Surgery*, 38(4), 604–609. doi:10.1111/j.1524-4725.2011.02223.x
- Al-Qattan, M. M. (2000). Surgical management of post-burn skin dyspigmentation of the upper limb. *Burns*, 26(6), 581–586. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10869832>
- American Academy of Dermatology. (1999). *Position statement on dispensing*.
- Arora, P., Sarkar, R., Garg, V. K., & Arya, L. (2012). Lasers for treatment of melasma and post-inflammatory hyperpigmentation. *Journal of Cutaneous and Aesthetic Surgery*, 5(2), 93–103. doi:10.4103/0974-2077.99436
- Baliña, L. M., & Graupe, K. (1991). The treatment of melasma. 20% azelaic acid versus 4% hydroquinone cream. *International Journal of Dermatology*, 30(12), 893–895. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1816137>
- Baumann, L. (2012). Ethics in cosmetic dermatology. *Clinics in Dermatology*, 30(5), 522–527. doi:10.1016/j.clindermatol.2011.06.023
- Boonchai, W., Desomchoke, R., & Lamtharachai, P. (2011). Trend of contact allergy to cosmetic ingredients in Thais over a period of 10 years. *Contact Dermatitis*, 65(6), 311–316.
- Britton, W. J., & Lockwood, D. N. (2004). Leprosy. *Lancet*, 363(9416), 1209–1219. doi:10.1016/S0140-6736(04)15952-7
- Cardinali, G., Kovacs, D., & Picardo, M. (2012). Mechanisms underlying post-inflammatory hyperpigmentation: Lessons from solar lentigo. *Annales de Dermatologie et de Vénérologie*, 139(Suppl. 4), S148–S152. doi:10.1016/S0151-9638(12)70127-8
- Chan, R., Park, K. C., Lee, M. H., Lee, E. S., Chang, S. E., Leow, Y. H., ... Verallo-Rowell, V. (2008). A randomized controlled trial of the efficacy and safety of a fixed triple combination (fluocinolone acetonide 0.01%, hydroquinone 4%, tretinoin 0.05%) compared with hydroquinone 4% cream in Asian patients with moderate to severe melasma. *The British Journal of Dermatology*, 159(3), 697–703. doi:10.1111/j.1365-2133.2008.08717.x
- Chow, M., & Jacob, S. E. (2015). The language of dermatology: The eponyms and inheritance of the genetic disorders of pigmentation. *Journal of the Dermatology Nurses' Association*, accepted for publication.
- Costin, G. E., & Hearing, V. J. (2007). Human skin pigmentation: Melanocytes modulate skin color in response to stress. *EASEB Journal*, 21(4), 976–994. doi:10.1096/fj.06-6649rev
- Crespo-Erchiga, V., & Florencio, V. D. (2006). Malassezia yeasts and pityriasis versicolor. *Current Opinion in Infectious Diseases*, 19(2), 139–147. doi:10.1097/01.qco.0000216624.21069.61
- Cristaudo, A., D'Illo, S., Gallinella, B., Mosca, A., Majorani, C., Violante, N., ... Petrucci, F. (2013). Use of potentially harmful skin-lightening products among immigrant women in Rome, Italy: A pilot. 226(3), 200–206.
- Dereure, O. (2001). Drug-induced skin pigmentation. Epidemiology, diagnosis and treatment. *American Journal of Clinical Dermatology*, 2(4), 253–262. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11705252>
- Dogra, S., Kanwar, A. J., & Parsad, D. (2002). Adapalene in the treatment of melasma: A preliminary report. *The Journal of Dermatology*, 29(8), 539–540. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12227492>
- Draelos, Z. D. (2007). Skin lightening preparations and the hydroquinone controversy. *Dermatologic Therapy*, 20(5), 308–313. doi:10.1111/j.1529-8019.2007.00144.x
- Eleftheriadou, V. (2013). Future horizons in vitiligo research: Focusing on the recommendations of the Cochrane systematic review "Interventions for vitiligo" 2010. *The British Journal of Dermatology*, 169(Suppl. 3), 67–70. doi:10.1111/bjd.12530
- Eleftheriadou, V., Thomas, K., Ravenscroft, J., Whitton, M., Batchelor, J., & Williams, H. (2014). Feasibility, double-blind, randomised, placebo-controlled, multi-centre trial of hand-held NB-UVB phototherapy for the treatment of vitiligo at home (HI-Light trial: Home Intervention of Light therapy). *Trials*, 15, 51. doi:10.1186/1745-6215-15-51
- Forste, G., Petrucci, F., & Bocca, B. (2008). Metal allergens of growing significance: Epidemiology, immunotoxicology, strategies for testing and prevention. *Inflammation & Allergy Drug Targets*, 7(3), 145–162.
- Gambichler, T., Krämer, H. J., Boms, S., Skrygan, M., Tomi, N. S., Altmeyer, P., & Maysner, P. (2007). Quantification of ultraviolet protective effects of pycnogenol in humans. *Archives of Dermatological Research*, 299(10), 517–520. doi:10.1007/s00403-007-0793-2
- Gilchrest, B. A., Blog, F. B., & Szabo, G. (1979). Effects of aging and chronic sun exposure on melanocytes in human skin. *The Journal of Investigative Dermatology*, 73(2), 141–143. doi:10.1111/1523-1747.ep12581580
- Gillbro, J. M., & Olsson, M. J. (2011). The melanogenesis and mechanisms of skin-lightening agents—Existing and new approaches. *International Journal of Cosmetic Science*, 33(3), 210–221. doi:10.1111/j.1468-2494.2010.00616.x
- Global leprosy update, 2013; reducing disease burden. (2014). *Relevé Épidémiologique Hebdomadaire (Weekly Epidemiological Record)*, 89(36), 389–400. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/25202781>



- Grimes, P. E. (1995). Melasma. Etiologic and therapeutic considerations. *Archives of Dermatology*, 131(12), 1453–1457. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7492140>
- Grimes, P. E., Bhawan, J., Guevara, I. L., Colón, L. E., Johnson, L. A., Gottschalk, R. W., & Pandya, A. G. (2010). Continuous therapy followed by a maintenance therapy regimen with a triple combination cream for melasma. *Journal of the American Academy of Dermatology*, 62(6), 962–967. doi:10.1016/j.jaad.2009.06.067
- Haake, A., & Holbrook, K. (1999a). The structure and development of skin. In T. B. Freedberg, I. M. Eisen, A. Z. Wolff, K. Austen, K. F. Goldsmith, L. A. Katz, & S. I. Fitzpatrick (Eds.), *Fitzpatrick's dermatology in general medicine* (pp. 70–114). New York, NY: McGraw-Hill.
- Halder, R. M., Nandedkar, M. A., & Neal, K. W. (2003). Pigmentary disorders in ethnic skin. *Dermatologic Clinics*, 21(4), 617–628. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14717403>
- Handel, A. C., Miot, L. D., & Miot, H. A. (2014). Melasma: A clinical and epidemiological review. *Anais Brasileiros de Dermatologia*, 89(5), 771–782. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4155956&tool=pmcentrez&rendertype=abstract>
- Hearing, V. J. (2000). The melanosome: The perfect model for cellular responses to the environment. *Pigment Cell Research*, 13(Suppl. 8), 23–34. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11041354>
- Hendrix, J. D. Jr., & Greer, K. E. (1992). Cutaneous hyperpigmentation caused by systemic drugs. *International Journal of Dermatology*, 31(7), 458–466. doi:10.1111/j.1365-4362.1992.tb02689.x
- Ho, S. G., Chan, N. P., Yeung, C. K., Shek, S. Y., Kono, T., & Chan, H. H. (2012). A retrospective analysis of the management of freckles and lentiginos using four different pigment lasers on Asian skin. *Journal of Cosmetic and Laser Therapy*, 14(2), 74–80. doi:10.3109/14764172.2012.670707
- Hughes, B. R. (1987). Melasma occurring in twin sisters. *Journal of the American Academy of Dermatology*, 17(5, Pt. 1), 841. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3680660>
- Hurwitz, E. (2006). *Hurwitz clinical pediatric dermatology: A textbook of skin disorders of childhood and adolescence*. Philadelphia, PA: Elsevier Health Sciences.
- Joshi, P. C., Carraro, C., & Pathak, M. A. (1987). Involvement of reactive oxygen species in the oxidation of tyrosine and dopa to melanin and in skin tanning. *Biochemical and Biophysical Research Communications*, 142(1), 265–274. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3101689>
- Katoulis, A. C., Stavrianeas, N. G., Georgala, S., Katsarou-Katsari, A., Koumantaki-Mathioudaki, E., Antoniou, C., & Stratigos, J. D. (1999). Familial cases of poikiloderma of Civatte: Genetic implications in its pathogenesis? *Clinical and Experimental Dermatology*, 24(5), 385–387. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10564328>
- Katoulis, A. C., Stavrianeas, N. G., Katsarou, A., Antoniou, C., Georgala, S., Rigopoulos, D., ... Katsambas, A. D. (2002). Evaluation of the role of contact sensitization and photosensitivity in the pathogenesis of poikiloderma of Civatte. *The British Journal of Dermatology*, 147(3), 493–497. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11207589>
- Kauvar, A. N. (2012). Successful treatment of melasma using a combination of microdermabrasion and Q-switched Nd:YAG lasers. *Lasers in Surgery and Medicine*, 44(2), 117–124. doi:10.1002/lsm.21156
- Kim, D., & Lockey, R. (2010). Dermatology for the allergist. *The World Allergy Organization Journal*, 3(6), 202–215. doi:10.1097/WOX.0b013e3181e2eb2e
- Lee, H. M., Haw, S., Kim, J. K., Chang, S. E., & Lee, M. W. (2013). Split-face study using a 1,927-nm thulium fiber fractional laser to treat photoaging and melasma in Asian skin. *Dermatologic Surgery*, 39(6), 879–888. doi:10.1111/dsu.12176
- Levantine, A., & Almeyda, J. (1973). Drug induced changes in pigmentation. *The British Journal of Dermatology*, 89(1), 105–112. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/4132858>
- Lin, R. L., & Janniger, C. K. (2005). Pityriasis alba. *Cutis*, 76(1), 21–24. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16144284>
- Makino, E. T., Mehta, R. C., Garruto, J., Gotz, V., Sigler, M. L., & Herndon, J. H. (2013). Clinical efficacy and safety of a multimodality skin brightener composition compared with 4% hydroquinone. *Journal of Drugs in Dermatology*, 12(3), s21–s26. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23545929>
- Molokhia, M. M., & Portmoy, B. (1973). Trace elements and skin pigmentation. *The British Journal of Dermatology*, 89(2), 207–209. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/4199347>
- Mulekar, S. V., & Isedeh, P. (2013). Surgical interventions for vitiligo: An evidence-based review. *The British Journal of Dermatology*, 169(Suppl. 3), 57–66. doi:10.1111/bjd.12532
- Nenoff, P., Krüger, C., Ginter-Hanselmayer, G., & Tietz, H. J. (2014). Mycology—An update. Part 1: Dermatofungal diseases: Causative agents, epidemiology and pathogenesis. *Journal der Deutschen Dermatologischen Gesellschaft (Journal of the German Society of Dermatology)*, 12(3), 188–209. doi:10.1111/ddg.12245
- Nicolaidou, E., & Katsambas, A. D. (2014). Pigmentation disorders: hyperpigmentation and hypopigmentation. *Clinics in Dermatology*, 32(1), 66–72. doi:10.1016/j.clindermatol.2013.05.026
- Ong, M. W., & Bashir, S. J. (2012). Fractional laser resurfacing for acne scars: A review. *The British Journal of Dermatology*, 166(6), 1160–1169. doi:10.1111/j.1365-2133.2012.10870.x
- Ortonne, J. (2008). Vitiligo and other disorders of hypopigmentation. In R. R. Bologna & J. Jorizzo (Eds.), *Dermatology* (2nd ed.). Oxford, UK: Elsevier Limited.
- Ortonne, J. P., Arellano, I., Berneburg, M., Cestari, T., Chan, H., Grimes, P., ... Westerhof, W. (2009). A global survey of the role of ultraviolet radiation and hormonal influences in the development of melasma. *Journal of the European Academy of Dermatology and Venereology*, 23(11), 1254–1262. doi:10.1111/j.1468-3083.2009.03295.x
- Ortonne, J. P., Pandya, A. G., Lui, H., & Hexsel, D. (2006). Treatment of solar lentigines. *Journal of the American Academy of Dermatology*, 54(5 Suppl. 2), S262–S271. doi:10.1016/j.jaad.2005.12.043
- Rigopoulos, D., Gregoriou, S., Charissi, C., Kontochristopoulos, G., Kalogeromitros, D., & Georgala, S. (2006). Tacrolimus ointment 0.1% in pityriasis alba: An open-label, randomized, placebo-controlled study. *The British Journal of Dermatology*, 155(1), 152–155. doi:10.1111/j.1365-2133.2006.07181.x
- Rivas, S., & Pandya, A. G. (2013). Treatment of melasma with topical agents, peels and lasers: An evidence-based review. *American Journal of Clinical Dermatology*, 14(5), 359–376. doi:10.1007/s40257-013-0038-4
- Sampogna, F., Raskovic, D., Guerra, L., Pedicelli, C., Tabolli, S., Leoni, L., ... Abeni, D. (2008). Identification of categories at risk for high quality of life impairment in patients with vitiligo. *The British Journal of Dermatology*, 159(2), 351–359. doi:10.1111/j.1365-2133.2008.08678.x
- Serrano, G., Pujol, C., Cuadra, J., Gallo, S., & Aliaga, A. (1989). Riehl's melanosis: Pigmented contact dermatitis caused by fragrances. *Journal of the American Academy of Dermatology*, 21(5, Pt. 2), 1057–1060. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2808836>
- Sheth, V. M., & Pandya, A. G. (2011). Melasma: A comprehensive update: Part II. *Journal of the American Academy of Dermatology*, 65(4), 699–714. doi:10.1016/j.jaad.2011.06.001
- Taieb, A., Alomar, A., Böhm, M., Dell'anna, M. L., De Pase, A., Eleftheriadou, V., ... Union Européenne des Médecins Spécialistes (UEMS). (2013). Guidelines for the management of vitiligo: The European Dermatology Forum consensus. *The British Journal of Dermatology*, 168(1), 5–19. doi:10.1111/j.1365-2133.2012.11197.x
- Taylor, C. R., & Anderson, R. R. (1994). Ineffective treatment of refractory melasma and postinflammatory hyperpigmentation by Q-switched ruby laser. *The Journal of Dermatologic Surgery and Oncology*, 20(9), 592–597. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8089359>
- Tierney, E. P., & Hanke, C. W. (2009). Treatment of Poikiloderma of Civatte with ablative fractional laser resurfacing: Prospective study and review of the literature. *Journal of Drugs in Dermatology*, 8(6), 527–534. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19537378>
- Torreló, A., Zaballos, P., Colmenero, I., Mediero, I. G., de Prada, I., & Zambrano, A. (2005). Erythema dyschromicum perstans in children: A report of 14 cases. *Journal of the European Academy of Dermatology and Venereology*, 19(4), 422–426. doi:10.1111/j.1468-3083.2005.01203.x
- Ustianowski, A. P., & Lockwood, D. N. (2003). Leprosy: Current diagnostic and treatment approaches. *Current Opinion in Infectious Diseases*, 16(5), 421–427. doi:10.1097/01.qco.0000092813.64370.be
- Vashi, N. A., & Kundu, R. V. (2013). Facial hyperpigmentation: Causes and treatment. *The British Journal of Dermatology*, 169, 41–56. doi:10.1111/bjd.12536
- Vazirnia, A., & Jacob, S. E. (2014). Review ACDS' Allergen of the Year 2000–2015. *The Dermatologist*, 22(11).
- Werlinger, K. D., Guevara, I. L., González, C. M., Rincón, E. T., Caetano, R., Haley, R. W., & Pandya, A. G. (2007). Prevalence of self-diagnosed melasma among premenopausal Latino women in Dallas and Fort Worth, Tex. *Archives of Dermatology*, 143(3), 424–425. doi:10.1001/archderm.143.3.424
- Woolery-Lloyd, H. C., Keri, J., & Doig, S. (2013). Retinoids and azelaic acid to treat acne and hyperpigmentation in skin of color. *Journal of Drugs in Dermatology*, 12(4), 434–437. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23652891>
- Zaynoun, S., Rubeiz, N., & Kibbi, A. G. (2008). Ashy dermatoses—A critical review of the literature and a proposed simplified clinical classification. *International Journal of Dermatology*, 47(6), 542–544. doi:10.1111/j.1365-4632.2008.03625.x

For more than 36 additional continuing education articles related to neonatal care, go to [NursingCenter.com](http://NursingCenter.com) \CE.